

3. REMUZZI G, WEENING JJ: Albuminuria as early test for vascular disease. *Lancet* 365:556–557, 2005
4. RITZ E: Treating microalbuminuria—Cosmetic exercise with urine chemistry or effective reduction of clinical events? *J Am Soc Nephrol* 16:1159–1163, 2005
5. ASSELBERGS FW, DIERCKS GFH, HILLEGE HL, *et al*: Effects of fosinopril and pravastatin on cardiovascular events in subjects with microalbuminuria. *Circulation* 110:2809–2816, 2004
6. THE HEART OUTCOMES PREVENTION EVALUATION STUDY INVESTIGATORS: Effects of an angiotensin converting enzyme inhibitor, ramipril, on cardiovascular events in high risk patients. *N Engl J Med* 342:145–153, 2000
7. IBSEN H, WACHTELL K, OLSEN MH, *et al*: Does albuminuria predict cardiovascular outcome on treatment with losartan versus atenolol in hypertension with left ventricle hypertrophy? A LIFE substudy. *J Hypertens* 22:1805–1811, 2004
8. IBSEN H, OLSEN MH, WACHTELL K, *et al*: Reduction of albuminuria translates to reduction in cardiovascular events in hypertensive patients. *Hypertension* 45:198–202, 2005

Diaphragmed fenestrae in the glomerular endothelium versus nondiaphragmed fenestrae in the hepatic endothelium

To the Editor: In a recent issue of *Kidney International*, Ballermann [1] discussed the development of diaphragmed fenestrae in glomerular endothelial cells. Major similarities exist between the hepatic and the glomerular endothelium regarding the process of fenestrae formation; intriguingly, an analogous controversy about the origin of fenestrae exists in both the hepatology and nephrology fields [2].

Hepatic endothelial fenestrae are dynamic structures that act as a sieving barrier to control the extensive exchange of material between the blood and the liver parenchyma. These membrane-bound pores lack a diaphragm and a basal lamina is also absent (Fig. 1 A and B, arrows), in contrast to fenestrae described in the kidney, pancreas, and brain. Their biological relevance in various diseases has been widely acknowledged [3].

Evidence that VEGF-induced diaphragmed fenestrae are derived from fused caveolae has recently been accumulating. It has been postulated that a similar mechanism is used in the formation of hepatic fenestrae. Conversely, actin disruption in hepatic endothelial cells produces an increase in the number of fenestrae by using a distinct structure that appears to serve as a fenestrae-forming center (Fig. 1C) [4]. Furthermore, it is important to emphasize the contradictions in the data reported about the origin of fenestrae in kidney, pancreas, and in the liver. For example, some data either clearly or indirectly indicate a possible relationship between caveolae, vesiculo-

vacuolar organelles, and (diaphragmed) fenestrae; others suggest the opposite (reviewed in [2]). Therefore, until this controversy is settled, no firm conclusions can be drawn, and stating that fenestrae, whether diaphragmed or not, correspond to fused and interconnected caveolae should be viewed with caution and remains as an emerging research topic.

FILIP BRAET and LILIAN L. SOON
Sydney, Australia

Correspondence to Filip Braet, Australian Key Centre for Microscopy and Microanalysis, Electron Microscope Unit, Madsen Building F09, University of Sydney, Sydney, NSW 2006, Australia.
E-mail: filip.braet@emu.usyd.edu.au

REFERENCES

1. BALLERMANN B: Glomerular endothelial cell differentiation. *Kidney Int* 67:1668–1671, 2005
2. BRAET F: How molecular microscopy revealed new insights in the dynamics of hepatic endothelial fenestrae in the past decade. *Liver Int* 24:532–539, 2004
3. BRAET F, WISSE E: Structural and functional aspects of liver sinusoidal endothelial cell fenestrae: A review. *Comparative Hepatology* 1:1, 2002
4. BRAET F, SPECTOR I, DE ZANGER R, WISSE E: A novel structure involved in the formation of liver endothelial cell fenestrae revealed using the actin inhibitor misakinolide. *Proc Natl Acad Sci USA* 95:13635–13640, 1998

Reply from the Authors

Braet and Soon question the conclusions we have drawn in a recent review [1] about the relationship between caveolae and fenestrae in glomerular endothelium. They suggest that we were too hasty in stating that fenestrae do not represent fused caveolae.

There is little doubt about the dynamic nature of fenestrae. Work with liver sinusoidal endothelial cells is particularly interesting because these cells retain very nice sieve plate structures in culture which can change rapidly by alterations in actin microfilament dynamics [2]. Also, VEGF-A can induce endothelial fenestration. Conversely, interference with VEGF-A function in renal glomeruli, for instance, in preeclampsia [3] or with podocyte VEGF-A haploinsufficiency [4] leads to an absence of glomerular fenestrae. Finally, VEGF-A seems to induce caveolin-1-containing vesicles to fuse and to form transendothelial cell channels [5, 6]. Such circumstantial evidence has suggested that fenestrae represent fused caveolae.

However, in caveolin-1-deficient mice known to lack caveolae altogether [7], we found glomerular endothelium with fenestrae that were normal in size and density, even though no caveolin-1 and no caveolae were observed at all [8]. In normal rat glomeruli examined for caveolin-1 localization by immunogold electron microscopy, fenestrated portions of glomerular




Fig. 1. High magnification electron micrographs of rat hepatic endothelial fenestrae. (A) Transmission electron microscopy image of sectioned hepatic endothelial cells *in situ*, showing the fenestrated processes (arrow) and the histologic relationship with the numerous microvillous processes of the liver parenchymal cells situated in the space of Disse. (B) Scanning electron micrograph of the fenestrated cytoplasm of hepatic endothelial cells *in vitro*. Notice the fenestrae (arrow) on the cell surface. (C) Whole-mount transmission electron micrograph of the fenestrated cytoplasm (arrow) of a hepatic endothelial cell cultured on transmission electron microscopy grids, disclosing the process of fenestrae formation. Note the area of intermediated density from which rows of fenestrae with increasing diameter are radiating into the surrounding cytoplasm. Scale bars, 200 nm.

endothelium were completely devoid of caveolin-1, similar to the report by Esser et al [9] for the choroid plexus endothelium. Now, our experiments leave open the possibility that fenestrae in other vascular beds could represent fused caveolae, and that caveolar precursor vesicles lacking caveolin-1 could fuse to form fenestrae. But if caveolae are defined as specialized plasma membrane invaginations stabilized by caveolins, then our findings in caveolin-deficient mice seem to prove that fenestrae in glomerular endothelium cannot represent fused caveolae.

BARBARA J. BALLERMANN
Edmonton, Alberta, Canada

Correspondence to Barbara J. Ballermann, Director of Nephrology,
University of Alberta, Edmonton, Alberta, Canada.
E-mail: barbara.ballermann@ualberta.ca

REFERENCES

1. BALLERMANN BJ: Glomerular endothelial cell differentiation. *Kidney Int* 67:1668–1671, 2005
2. BRAET F, WISSE E: Structural and functional aspects of liver sinusoidal endothelial cell fenestrae: A review. *Comp Hepatol* 1:1, 2002
3. MAYNARD SE, MIN JY, MERCHAN J, et al: Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest* 111:649–658, 2003
4. EREMINA V, SOOD M, HAIGH J, et al: Glomerular-specific alterations

- of VEGF-A expression lead to distinct congenital and acquired renal diseases. *J Clin Invest* 111:707–716, 2003
5. VASILE E, QU H, DVORAK HF, et al: Caveolae and vesiculo-vacuolar organelles in bovine capillary endothelial cells cultured with VPF/VEGF on floating Matrigel-collagen gels. *J Histochem Cytochem* 47:159–167, 1999
6. YOKOMORI H, ODA M, YOSHIMURA K, et al: Vascular endothelial growth factor increases fenestral permeability in hepatic sinusoidal endothelial cells. *Liver Int* 23:467–475, 2003
7. RAZANI B, ENGELMAN JA, WANG XB, et al: Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *J Biol Chem* 276:38121–38138, 2001
8. SORENNSSON J, FIERLBECK W, HEIDER T, et al: Glomerular endothelial fenestrae in vivo are not formed from caveolae. *J Am Soc Nephrol* 13:2639–2647, 2002
9. ESSER S, WOLBURG K, WOLBURG H, et al: Vascular endothelial growth factor induces endothelial fenestrations in vitro. *J Cell Biol* 140:947–959, 1998

Immunosuppressive and calcimimetic drug-drug interactions

To the Editor: Drug-drug interactions (DDIs) are an important cause of adverse drug reactions. It has been estimated that approximately 5% of prescribing errors [1] or of adverse drug reactions [2] are caused by DDIs in